

Plant Tissue Sampling

Daniel Geisseler, Patricia Lazicki and William R. Horwath

Background

- Plant analyses are useful to diagnose nutritional problems and to monitor the fertilization program. Tissue testing is most effective when used together with soil testing ^[13].
- Nutrient concentrations change as plants grow and also differ between plant parts ^[6]. It is therefore important to sample specific plant parts at a particular growth stage (Table 1). For information on optimal nutrient

concentrations at different growth stages, see the fertilization guidelines for the different crops (http://apps.cdfa.ca.gov/frep/docs/Intro.html)

 Archiving the results from the analyses allows tracking changes in the same field over time ^[6]. Plant analyses together with soil analyses and nutrient budgets allow evaluating the fertilization program on the long term ^[4].

General Sampling Instructions

- When plant development differs within a field, the field should be divided into different management areas with similar characteristics and a sample from each area should be taken. To facilitate interpretation, fields are best divided into the same areas as is done for soil samples.
- Randomly select plants throughout the field or management area and sample the correct plant parts ^[4, 13]. For plant parts and number of plant parts to sample, see Table 1.
- Collect the specific plant parts and place them into a clean paper bag ^[13]. Do not use plastic bags to avoid decay of samples. Do not use metal containers, because they may contaminate the samples and affect micronutrient results ^[4, 14].
- Do not collect samples during the hottest part of the day, particularly in summer ^[14].
- Do not take samples from dead, diseased, insect damaged, or mechanically injured plants ^[6, 13]. Also avoid plants from unusual areas in the field, including border areas and

places where plants are under water stress or where nutrient availability is atypical ^[6].

- Dust or soil covered plant parts should also be avoided, especially when the samples are used for micronutrient analysis ^[6].
- Small amounts of dust can be removed by gently brushing the samples with a soft brush ^[13]. Alternatively, the samples may be cleaned with a damp cloth, but should not be rinsed or washed to prevent leaching of nutrients from the sample ^[13, 14].
- Deliver the samples immediately to the lab or use a one-day delivery service ^[13, 14]. If immediate delivery is not possible, air-dry the samples in the shade by placing the open bag in a clean, dust-free area ^[3, 4, 14]. Mix the samples frequently to avoid decay.
- Clearly label the bag, and provide the information required by the test lab ^[4].
- Follow the laboratory instructions for packaging and shipping.

• To determine the cause of visual symptoms or a suspected deficiency in one part of the field, two samples may be taken; one from the plants of interest, the other from adjoining normal plants in the same field or management area ^[6].

Table 1: Sampling procedure for major field crops

| Crop | Growth stage | Plant part to sample | Number of plants to sample |
|------------------------|---|--|--|
| Field Crops | | | |
| Alfalfa | 10% bloom | stems in the middle third of the plant | 40-60 stems from at least 30 plants |
| Dry Beans | Early growth | Petiole of fourth leaf from the growing tip | 40 |
| | Pre-bloom | Petiole of fourth leaf from the growing tip | 40 |
| | Late bloom | Petiole of fourth leaf from the growing tip | 40 |
| Corn | Early season (6-16 inches) | Whole plant | 20-30 |
| | Midgrowth (3-6 feet) | First fully developed leaf; third leaf from top | 15-25 |
| | Tasseling | Leaf opposite and below primary ear | 15-25 |
| | Silking | Leaf opposite and below primary ear | 15-25 |
| Cotton | Early squaring to late season | Third to fifth petiole from the terminal on the main stem | 30-40 |
| Rice | Early stages | Most recently fully expanded leaf (Y-leaf) | 50 |
| | Later stages | Most recently fully expanded leaf (Y-leaf) | 30-60 |
| Safflower | Prebloom | Stem, middle section | 40 |
| | First bud visible | Recently matured mid-stem leaves | 40 |
| Sunflower | Just before floret emergence | Most recently matured leaf (3rd or 4th from the terminal bud | 20-30 |
| Processing Tomatoes | First bloom to 10% of fruits showing red color | Fourth leaf from the growing tip | 40 |
| Wheat and barley | 3-4 leaf | Whole plant | 50-100 |
| | Tillering | Top 3-4 leaves | 50-100 |
| | Jointing | Top 3-4 leaves | 50-100 |
| | Booting | Top 3-4 leaves | 50-100 |
| | Early heading (hard red wheat N only) | Flag leaf | 50-100 |

Sources: Alfalfa^[9,12], dry beans^[2], corn^[6], cotton^[1,11], rice^[10], safflower^[2,7], sunflower^[2,4,7], tomatoes^[7,8], wheat and barley^[6]

| Crop | Growth stage | Plant part to sample | Number of plants to sample |
|-------------|---------------------------------|---|-------------------------------|
| Vegetables | | | |
| Broccoli | First buds to heading | Recently matured leaf, typically 3-4 nodes down from the growing point | 20-60 |
| Carrot | Midgrowth (>4 inches high) | Most recently matured leaf or petiole | 20-30 |
| Cauliflower | Head initiation | Recently matured leaf, typically 3-4 nodes down from the growing point | 20 |
| | Preharvest | Recently matured leaf, typically 3-4 nodes down from the growing point | 20 |
| Celery | Mid-growth | Most recently matured leaf or petiole | 20 |
| | Preharvest | Most recently matured leaf or petiole | 20 |
| Melon | Early flower | Most recently matured leaf or petiole, typically 6th from the growing tip | 20-30 |
| | Early fruit set/bulking | Most recently matured leaf or petiole, typically 6th from the growing tip | 20-30 |
| | First harvest | Most recently matured leaf or petiole, typically 6th from the growing tip | 20-30 |
| Onion | Early season | Tallest leaf | 20-30 |
| | Midseason | Tallest leaf | 20-30 |
| | Late season | Tallest leaf | 20-30 |
| Potato | Early season | Petiole of fourth leaf from the growing tip | 40 |
| | Midseason | Petiole of fourth leaf from the growing tip | 40 |
| | Late season | Petiole of fourth leaf from the growing tip | 40 |
| Lettuce | Early heading to pre-harvest | Youngest wrapper leaf | 20-60 |
| Berries | | | |
| Strawberry | Preharvest | Young mature leaves | 30-40 |
| | Main harvest | Young mature leaves | 30-40 |

 Table 2: Sampling procedure for major vegetable and berry crops

Sources: Broccoli, cauliflower, celery and lettuce ^[5, 6], carrot ^[2,6], melon ^[2,7], onion ^[8], potato ^[7,16], strawberry ^[15]

References

- Bassett, D.M., MacKenzie, A.J., 1976. Plant analysis as a guide to cotton fertilization. In: Reisenauer, H.M. (Ed.). Soil and Plant-Tissue Testing in California. University of California Cooperative Extension Bulletin 1879. pp. 16-17.
- California Plant Health Association, 2002. Western Fertilizer Handbook 9th edition. Interstate Publishers, Inc.
- Fernandez, F.G., Hoeft, R.G., 2012. Managing soil pH and crop nutrients. University of Illinois Extension -Illinois Agronomy Handbook. Available online at: <u>http://extension.cropsci.illinois.edu/handbook/pdfs/chapt er08.pdf</u>
- Flynn, R., Ball, S.T., Baker, R.D., 1999. Sampling for plant tissue analysis. New Mexico State University Cooperative Extension Service. Available online at: <u>http://aces.nmsu.edu/pubs/_a/A123.pdf</u>
- Hartz, T.K., 2007. Efficient nitrogen management for cool-season vegetables. Available online at: <u>http://vric.ucdavis.edu/pdf/fertilization_EfficientNitrogen</u> <u>ManagementforCoolSeasonVegetable2007.pdf</u>
- Jones Jr., J.B., 1998. Field sampling procedures for conducting a plant analysis. In: Kalra, Y.P. (Ed.). Handbook of Reference Methods for Plant Analysis. CRC Press, Boca Raton. pp. 25-35.
- Lorenz, O.A., Tyler, K.B., 1976. Plant tissue analysis of vegetable crops. In: Reisenauer, H.M. (Ed.). Soil and Plant-Tissue Testing in California. University of California Cooperative Extension Bulletin 1879. pp. 24-29.
- Maynard, D.N., Hochmuth, G.J., 2007. Knott's Handbook for Vegetable Growers. John Wiley & Sons, Inc., Hoboken, NJ.
- Meyer, R.D., Marcum, D.B., Schmierer, J.L., 1998. Do I need fertilizer? - Ask the plant! Proceedings of the 28th

California/Nevada Alfalfa Symposium.Available online at:

http://alfalfa.ucdavis.edu/+symposium/proceedings/199 8/98-159.pdf

- Mikkelsen, D.S., 1976. Diagnostic plant analysis for rice. In: Reisenauer, H.M. (Ed.). Soil and Plant-Tissue Testing in California. University of California Cooperative Extension Bulletin 1879. pp. 30-31.
- Reddy, K.R., Hodges, H.F., Varco, J., 2000. Potassium nutrition of cotton. Mississippi Agricultural and Forestry Experiment Station Bulletin 1049. Available online at <u>http://msucares.com/pubs/bulletins/b1094.pdf</u>
- Sallee, W.R., Ulrich, A., Martin, W.E., Krantz, B.A., 1959. High phosphorus for alfalfa: Plant analysis used to evaluate phosphorus status of alfalfa fields as guide to fertilizing for better yields and returns. California Agriculture 13(8), 7-8.
- Schwab, G.J., Lee, C.D., Pearce, R., 2007 Sampling plant tissue for nutrient analysis. University of Kentucky Cooperative Extension Service. Available online at: <u>http://www.ca.uky.edu/agc/pubs/agr/agr92/agr92.pdf</u>
- Thom, W.O., Brown, J.R., Plank, C.O., 1991. Sampling for corn plant tissue analysis. Iowa State University Extension. Available online at: <u>http://www.extension.iastate.edu/Publications/NCH15.p</u> <u>df</u>
- Ulrich, A., 1976. Plant tissue analysis as a guide in fertilizing crops. In: Reisenauer, H.M. (Ed.). Soil and Plant-Tissue Testing in California. University of California Cooperative Extension Bulletin 1879. pp. 6-8
- Voss, R.E., 2004. Reducing fertilizer needs of potato with new varieties and clonal strains of existing varieties. FREP final report. Available online at: <u>http://www.cdfa.ca.gov/is/ffldrs/frep/pdfs/completedproj</u> <u>ects/00-0514Voss2004.pdf</u>

Daniel Geisseler is an Extension Specialist in the Department of Land, Air and Water Resources at the University of California, Davis.

Patricia Lazicki is an Assistant Specialist in the Department of Land, Air and Water Resources at the University of California, Davis.

William R. Horwath is professor of Soils and Biogeochemistry in the Department of Land, Air and Water Resources and the James G. Boswell Endowed Chair in Soil Science at the University of California, Davis.

The document has been prepared within the project "Assessment of Plant Fertility and Fertilizer Requirements for Agricultural Crops in California", funded by the California Department of Food and Agriculture Fertilizer Research and Education Program (FREP).

This document is available online at https://apps1.cdfa.ca.gov/FertilizerResearch/docs/Plant_Tissue_Sampling.pdf

Last update: January, 2017